

Total No. of printed pages =03

SUBJECT CODE: INT052102

Roll No. of candidate

--	--	--	--	--	--	--	--	--	--

2017
End Semester B.Sc.(IT) Examination
1st Semester
MATHEMATICS-I

Full Marks- 70

Pass Marks : 21

Time- 3 hours

The figures in the margin indicate full marks for the questions.

PART – A

Q1. Answer all questions:

[16 × 1 = 16]

- Give an example of a bounded above sequence.
- What are the limit points of the sequence $\{(-1)^n\}$?
- Under what condition a monotonic sequence becomes convergent?
- What is alternating series?
- Define a group.
- Give an example of a ring.
- State Lagrange's theorem on the order of a subgroup of a finite group.
- Give an example of a field.
- If α, β, γ are roots of the equation $x^3 - 7x^2 + 36 = 0$, then find the value of $\alpha + \beta + \gamma$.
- If two roots of an cubic equation are 2 and $1 + i$, then find the third root of the equation.
- State Descarte's rule of sign for roots of an equation.
- Write the necessary condition for maxima and minima of a function $f(x)$ at point $= c$.
- Find modulus of the complex number $= \frac{1+i}{1-i}$.
- Write the complex number $z = \sqrt{3} - i$ in polar form.
- If $z = \cos\theta + i\sin\theta$, then find $z^3 + \frac{1}{z^3}$.
- If z is a complex number, then show that $\cos(iz) = \cosh(z)$.

P.T.O.

PART – B

Q2. Answer all questions:

[4 × 3.5 = 14]

- a) Check the convergence of the sequence $\{n + (-1)^n n\}$.
- b) Give an example to show that union of two subgroups is not necessarily a subgroup.
- c) If α, β, γ are roots of the equation $x^3 - px^2 + qx - r = 0$, then find the value of $\sum \alpha^2$.
- d) Evaluate the Gregory series $\left(\frac{2}{3} + \frac{1}{7}\right) - \frac{1}{3}\left(\frac{2}{3^3} + \frac{1}{7^3}\right) + \frac{1}{5}\left(\frac{2}{3^5} + \frac{1}{7^5}\right) - \dots \infty$.

PART – C

(All questions are compulsory)

Q.3.

[5 + 5 = 10]

- a) Show that a sequence cannot converge to more than one limit.
- b) Show that the series $1 - \frac{1}{2^3} + \frac{1}{3^3} - \dots$ is absolutely convergent.

OR

- a) Show that the sequence $\{S_n\}$ where $S_n = \frac{1}{n+1} + \frac{1}{n+2} + \dots + \frac{1}{n+n}$ is convergent.
- b) Test the convergency of $\sum \frac{n^2-1}{n^2+1} x^n, x > 0$, using D' Alembert's ratio test.

Q4.

[5 + 5 = 10]

- a) Use De Moivre's theorem to prove that $\tan 5\theta = \frac{5 \tan \theta - 10 \tan^3 \theta + \tan^5 \theta}{1 - 10 \tan^2 \theta + 5 \tan^4 \theta}$.
- b) Prove that $\frac{(\cos \theta + i \sin \theta)^4}{(\cos \theta + \sin \theta)^5} = \sin 9\theta - i \cos 9\theta$ and find all the values of $(1 - i\sqrt{3})^{1/4}$.

OR

- a) Expand $\sin^3 x$ in power of x .
- b) If x, y are real numbers, then show that
 - i) $\sinh(x + iy) = \sinh x \cos y + i \cosh x \sin y$
 - ii) $\cos(x + y) = \cosh x \cosh y + \sinh x \sinh y$

Q5.

[5 + 5 = 10]

- a) If $f = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix}, g = \begin{pmatrix} 2 & 1 & 3 \\ 2 & 1 & 3 \end{pmatrix}$, show that $fg \neq gf$.
- b) Prove that every cyclic group is abelian.

OR

- a) Show that the mapping $f: R \rightarrow R^+$ given by $f(a) = 2^a, a \in R$, where R is the group of real numbers under addition and R^+ , the group of positive real numbers under multiplication is a homomorphism.
- b) If R is a ring, then show that $a0 = 0a = 0$, for every $a \in R$ and where 0 is the additive identity of the ring R .

P.T.O.

Q6.

[5 + 5 = 10]

- a) Solve the cubic equation by Cardon's method $x^3 - 27x + 54 = 0$.
- b) Solve the cubic equation $x^3 - 5x^2 - 2x + 24 = 0$, given that the product of two of the roots is 12.

OR

- a) Find for what values of x , the expression $x^3 - 9x^2 + 15x - 3$ is maximum and minimum. Also find its maximum and minimum values.
- b) Evaluate $\lim_{x \rightarrow 0} \frac{e^x - e^{-x} - 2x}{x - \sin x}$
